

Observation of a metallic superfluid in a numerical experiment

E. Smørgrav,¹ E. Babaev,^{2,1} J. Smiseth,¹ and A. Sudbø¹

¹*Department of Physics, Norwegian University of Science and Technology, N-7491 Trondheim, Norway*

²*Laboratory of Atomic and Solid State Physics, Cornell University, Ithaca, NY 14853-2501, USA*

We report the observation, in Monte Carlo simulations, of a novel type of quantum ordered state: *the metallic superfluid*. The metallic superfluid features ohmic resistance to counter-flows of protons and electrons, while featuring dissipationless co-flows of electrons and protons. One of the candidates for a physical realization of this remarkable state of matter is hydrogen or its isotopes under high compression. This adds another potential candidate to the presently known quantum dissipationless states, namely superconductors, superfluid liquids and vapours, and supersolids.

PACS numbers: 71.10.Hf, 74.10.+v, 74.90.+n, 11.15.Ha

At low temperatures, fluids become dominated by the wavelike nature of their constituent particles when the thermal de Broglie wavelength exceeds the interparticle separation. Such quantum fluids usually feature superconductivity or superfluidity, which however may be destroyed by topological line-defects (vortices) threading the entire system. Vortices may be induced by a magnetic field or rotation¹, or by thermally excited *transverse* phase fluctuations of the macroscopic wavefunction of superconductors and superfluids^{2,3,4}. In a system of rapidly growing interest, a two-component superconductor⁵, vortices yield dramatic physical consequences. Here we report the first observations, in a numerical experiment, of a novel type of quantum fluid originating in aggregate states of vortex matter. Increasing temperature, a composite vortex lattice melts into a composite vortex liquid, whence superconductivity is lost while superfluidity is retained, *yielding a metallic (ohmic) superfluid*. At higher temperature, another unusual transition occurs where the composite vortex liquid “ionizes” into a “plasma” of constituent vortices, destroying superfluidity.

Recently, there has been considerable interest in theories of superconductors with several superconducting components coupled by a magnetic field, but with no possibility of Josephson tunneling of one component into another. This is predicted to occur in a wide variety of physical systems, most notably in condensed matter such as hydrogenic atoms subjected to extreme pressure^{5,6,7,8,9,10} or effective theories of easy-plane quantum antiferromagnets¹¹. Renewed interest in the long sought liquid metallic hydrogen (LMH) is due to recent ab initio calculations¹² along with a breakthrough in synthesis of ultrahard artificial diamonds, essential for obtaining the required extreme pressures in anvil cells¹³. These facts, along with a recent measurement of an unusually low melting temperature of dense Na¹⁴, strongly hint at a realization of LMH in the not too distant future. Thus, understanding the superconducting properties of this system is important, since magnetic field experiments can be conducted in high pressure anvil cells, possibly confirming the realization of this novel state of matter in a terrestrial laboratory. It is commonly accepted that

LMH is abundant in the interior of Jupiter and Saturn and quite possibly also present in some of the known 200 extrasolar giant planets¹⁵. In these cases, however, LMH is conjectured to exist in the classical metallic regime at several thousand degrees Kelvin. In contrast, the state projected to exist in Refs. 6 is a ground state quantum fluid, and is the one we focus on in this Letter.

Realization of LMH could well constitute the next milestone in quantum fluids. It is projected to feature Cooper-pairs of both electrons and protons at low temperatures⁶. The resulting quantum fluid differs radically from previously known quantum fluids, in that its physical properties cannot be classified exclusively as a superconductor or a superfluid⁵. Remarkably, such a system features both superconductivity and superfluidity which appear as collective phenomena corresponding to co- and counter-flows of two species of Cooper-pairs, with a complicated interplay between them. Thus, LMH has been conjectured to sustain phase transitions connecting a superconducting *and* superfluid state to a metallic state featuring a superfluid mode⁵, or to a superconducting state with no superfluid mode.

The transition from a state featuring superconductivity *and* superfluidity to a state where superfluidity is lost and superconductivity is retained, has recently been observed in a large scale Monte Carlo (MC) simulation¹⁰. However, the remarkable possibility of a transition from a “composite” vortex lattice (superconducting state where vortex matter forms a solid) to a “composite vortex liquid” (metallic superfluid state of the system) along with a subsequent transition from a composite vortex liquid to “vortex line plasma”, *has thus far not been confirmed*. *It is the purpose of this Letter to report an observation of the metallic superfluid, not dealt with in simulations previously, in a numerical experiment.*

The superconducting phase of LMH is given by the Ginzburg-Landau model with two scalar fields $\Psi_0^{(1)}(\mathbf{r})$ and $\Psi_0^{(2)}(\mathbf{r})$ describing superconducting condensates of protons and electrons, respectively. It is defined via the energy density

$$\mathcal{H} = \sum_{\alpha=1}^2 \frac{|\mathbf{D}\Psi_0^{(\alpha)}(\mathbf{r})|^2}{2M^{(\alpha)}} + V(|\Psi_0^{(\alpha)}(\mathbf{r})|) + \frac{1}{2}(\nabla \times \mathbf{A}(\mathbf{r}))^2. (1)$$

Here, $M^{(1)}$ and $M^{(2)}$ are the masses of the condensates, the covariant derivative is given by $\mathbf{D} \equiv \nabla - ie\mathbf{A}(\mathbf{r})$, and $V(\{|\Psi_0^{(\alpha)}(\mathbf{r})|\})$ is the potential term. The particular form of $V(\{|\Psi_0^{(\alpha)}(\mathbf{r})|\})$ is not essential for the large-scale physics we address, but its dependence only on $|\Psi_0^{(\alpha)}|$ reflects the fact that Cooper-pairs of electrons cannot be converted into Cooper-pairs of protons, and vice versa. For the issues discussed in this paper, it suffices to work in the phase only approximation $\Psi_0^{(\alpha)}(\mathbf{r}) = |\Psi_0^{(\alpha)}| \exp[i\theta^{(\alpha)}(\mathbf{r})]$ where $|\Psi_0^{(\alpha)}|$ is treated as a constant¹⁰. In LMH, the two order parameters correspond to electronic and protonic Cooper-pairs. Moreover, Eq. (1) may be rewritten as follows^{5,7,8,10}

$$\mathcal{H} = \frac{1}{2\Psi^2} \left(|\psi^{(1)}|^2 \nabla \theta^{(1)} + |\psi^{(2)}|^2 \nabla \theta^{(2)} - e\Psi^2 \mathbf{A} \right)^2 \quad (2)$$

$$+ \frac{1}{4\Psi^2} |\psi^{(1)}|^2 |\psi^{(2)}|^2 \left(\nabla(\theta^{(1)} - \theta^{(2)}) \right)^2 + \frac{1}{2} (\nabla \times \mathbf{A})^2,$$

where $|\psi^{(\alpha)}|^2 = |\Psi_0^{(\alpha)}|^2 / M^{(\alpha)}$, and $\Psi^2 \equiv |\psi^{(1)}|^2 + |\psi^{(2)}|^2$. The neutral and charged modes, described by the second and first term in Eq. (2), are explicitly identified.

The topological objects of Eq. (1) are vortices of type-1 and type-2 defined by a 2π winding in $\theta^{(1)}$ and $\theta^{(2)}$, respectively. The interaction potential between these vortices is a superposition of a Coulomb potential and a Yukawa potential, arising out of the neutral and the charged mode, respectively. (For a derivation of the formulae for the vortex interaction, see Ref. 10). The energy of a vortex associated with $\pm 2\pi$ winding in $\theta^{(1)} - \theta^{(2)}$ is logarithmically divergent. As the neutral mode tends to lock $\nabla \theta^{(1)}$ to $\nabla \theta^{(2)}$, in order to minimize the second term in Eq. (2), the vortices of type-1 and type-2 pair up into a composite vortex for which $\nabla(\theta^{(1)} - \theta^{(2)}) = 0$. Therefore, a composite vortex is an object where a type-1 and a type-2 vortex are co-centered and co-directed in space, and the Coulomb part of the pair-potential exactly cancels. The screened potentials add, but the associated overall energy is finite^{5,8,9,10}. In the presence of an externally applied magnetic field, the ground state of the system is a lattice of co-centered vortices of type-1 and type-2, a *composite vortex lattice*.

We have performed MC simulations on Eq. (1) at finite temperature using local Metropolis updating on the fields $\theta^{(1)}(\mathbf{r})$, $\theta^{(2)}(\mathbf{r})$, and $A(\mathbf{r})$. The system size we have used is $L \times L \times L$, with $L = 120$. The coupling constants investigated are $|\psi^{(1)}|^2 = 0.5$, $|\psi^{(2)}|^2 = 1.0$ and $e = 1.0$ and the external magnetic field is $\mathbf{B} = \nabla \times \mathbf{A}(\mathbf{r}) = (0, 0, 2\pi f)$ with $f = 1/20$. Thus, there are 20 plaquettes in the xy -plane for each field-induced vortex. The external magnetic field is imposed by splitting $\mathbf{A}(\mathbf{r}) = \mathbf{A}_F(\mathbf{r}) + \mathbf{A}_0(\mathbf{r})$, where $\mathbf{A}_F$ is free to fluctuate subject to periodic boundary conditions, and $\mathbf{A}_0 = (0, 2\pi x f, 0)$ is kept fixed. We have chosen the amplitude ratios $|\psi^{(2)}|^2 / |\psi^{(1)}|^2 = 2.0$ for numerical convenience. We emphasize that the results in this paper are dictated by symmetry, and the physical picture we present will thus be representative also for LMH, where $|\psi^{(2)}|^2 / |\psi^{(1)}|^2 \sim 10^3$.

To find the lattice ordering of vortices we compute the planar structure function $S^{(\alpha)}(\mathbf{k}_\perp)$ of the local vorticity $\mathbf{n}^{(\alpha)}(\mathbf{r})$, defined by $\Delta \times [\Delta \theta^{(\alpha)} - e\mathbf{A}] = 2\pi \mathbf{n}^{(\alpha)}(\mathbf{r})$, given by $S^{(\alpha)}(\mathbf{k}_\perp) = \langle |\sum_{\mathbf{r}} n_z^{(\alpha)}(\mathbf{r}) e^{i\mathbf{k}_\perp \cdot \mathbf{r}_\perp}|^2 \rangle / (fL^3)^2$. Here, Δ_μ is the lattice difference operator, $\Delta_\mu \theta^{(\alpha)} - eA_\mu \in [0, 2\pi)$, $\mathbf{r}$ runs over the possible positions of the vortices, and $\mathbf{k}_\perp$ and $\mathbf{r}_\perp$ are perpendicular to $\mathbf{B}$. If vortices form a lattice, $S^{(\alpha)}(\mathbf{k}_\perp)$ will exhibit a six-fold symmetric Bragg structure, and feature a ring-structure in the vortex liquid phase.




FIG. 1: (Color online) Results of numerical experiments on a two-component vortex system, for $|\psi^{(1)}|^2 = 0.5$, $|\psi^{(2)}|^2 = 1.0$, and $e = 1.0$. Panel **a**: Structure functions for protonic $S^{(1)}(\mathbf{K})$ (red) and electronic $S^{(2)}(\mathbf{K})$ (blue) vortices, for $\mathbf{K} = (\pi/4, 2\pi/5)$. They drop to zero discontinuously at T_M where the system loses its superconducting properties, but retains superfluidity, as evidenced by a finite helicity modulus Υ_x (black line). Υ_x serves as an order parameter in the vortex liquid phase, dropping to zero at T_C . Panel **b**: N_{co} (green) is finite across the melting transition, but drops continuously to a small value at T_C , where it has a kink. At T_C , the specific heat C_V has an anomaly.

The MC results are given in Fig. 1, showing the structure function $S^{(1)}(\mathbf{K})$ (red) for protonic vortices, and $S^{(2)}(\mathbf{K})$ (blue) for electronic vortices, where $\mathbf{K} = (\pi/4, 2\pi/5)$ is a Bragg vector. In the low temperature regime both functions are finite, but decrease gradually as the temperature is increased. At T_M , $S^{(1)}(\mathbf{K})$ and $S^{(2)}(\mathbf{K})$ vanish at the same temperature even though the ratio of the bare stiffnesses of the condensates is 2.0. Moreover, both structure functions vanish discontinuously, the hallmark of a first order melting transition of the *composite vortex lattice*.

To probe the splitting of the co-centered vortices into constituent vortices, we compute the vortex co-centrality, defined as $N_{co} \equiv N_{co}^{(+)} - N_{co}^{(-)}$ where $N_{co}^{(\pm)} =$

$\sum_{\mathbf{r}} |n_z^{(2)}(\mathbf{r})| |\delta_{n_z^{(1)}(\mathbf{r}), \pm n_z^{(2)}(\mathbf{r})} / \sum_{\mathbf{r}} |n_z^{(2)}(\mathbf{r})|$, and $\delta_{i,j}$ is unity if $i = j$ and zero otherwise. Therefore, N_{co} is the fraction of type-2 vortex segments that are co-centered and co-linear with type-1 segments. We find that passing through the first-order melting transition renders N_{co} unaffected, whence we conclude that the observed transition is a melting of a *composite* vortex lattice.

In one-component type-II superconductors, melting of the vortex lattice amounts to a complete destruction of dissipationless currents⁴. To follow the fate of the superfluid mode of the two-component system in the vortex liquid state, we measure the ordering in $\gamma(\mathbf{r}) \equiv \theta^{(1)}(\mathbf{r}) - \theta^{(2)}(\mathbf{r})$. To probe the global phase coherence in this variable we consider the helicity modulus Υ_μ , equivalently the superfluid density, given by $\Upsilon_\mu = [\langle c \rangle - \eta \langle s^2 \rangle] / L^3$, where $c = \sum_{\mathbf{r}} \cos(\Delta_\mu \gamma(\mathbf{r}))$, $s = \sum_{\mathbf{r}} \sin(\Delta_\mu \gamma(\mathbf{r}))$, and $\eta = \beta |\psi^{(1)}|^2 |\psi^{(2)}|^2 / 2 \Psi^2$. When the composite vortex lattice melts, destroying superconductivity⁴, *the superfluid density remains unaffected*, cf. Fig. 1. *Hence, this transition separates a superconducting superfluid from a metallic superfluid state.*

Thus, we have identified a transition from a composite vortex lattice into a composite vortex liquid. Increasing the temperature further, we find a phase transition between the *composite vortex liquid* and the “*ionized vortices*” *plasma*, as evidenced by the vanishing of the superfluid density Υ_x , cf. Fig. 1. *We observe that this*

transition is accompanied by a pronounced anomaly in the specific heat C_V , indicating that this is a critical phenomenon. This is corroborated by the following physical argument. The liquid state of vortices in an ordinary superconductor is a state where translational symmetry is restored and superconductivity is lost⁴. In the *composite vortex liquid* state, every electronic vortex is accompanied by a protonic vortex performing only finite excursions away from the electronic vortex line. Therefore, for every plane slicing an electronic vortex in a direction perpendicular to the magnetic field, it is possible to identify a finite length closed contour which also encompasses an accompanying co-directed protonic vortex. Along such a contour, there is no nontrivial winding in the phase-difference γ . Therefore, melting of the composite vortex lattice into a composite vortex liquid does not restore the broken global $U(1)$ -symmetry associated with γ . On the other hand, in the vortex plasma state one cannot find a protonic vortex accompanying every electronic vortex, implying a disordering of γ . Hence, the associated global $U(1)$ -symmetry is restored during the “vortex ionization” transition taking place within the vortex liquid, and it is therefore in the $3Dxy$ universality class.

To gain further insight, we have also extracted vortices and visualized snapshots of configurations of vortex matter in small segments of the system. The results are shown in Fig. 2.

Panel **a** of Fig. 2 shows the side- and top-views of vortex matter at a small, but finite, temperature when two species of vortices perform only small excursion from each other. The insets show the structure function in momentum space of the protonic (thin red) and electronic (thick blue) vortices. Parallel and co-directed vortices interact with each other as positively and negatively charged strings, and the splitting is a temperature-induced fluctuation. Panel **b** shows snapshots of the vortex matter when the system is heated above the temperature of vortex lattice melting. The protonic (red) and electronic (blue) structure functions have developed ring-like structures, characteristic of a vortex-liquid in both the protonic and electronic sectors. While the vortices perform stronger excursions from each other, these excursions are still limited and one can always identify a red line attached to any given blue line. Thus, co-centricity of protonic (red) and electronic (blue) vortex lines is still largely intact. In such a configuration, a dissipationless electrical current cannot propagate in any direction. Quite remarkably, however, co-directed currents of protonic and electronic pairs can propagate through this system without dissipation, as evidenced by the measurements of the helicity modulus given in Fig. 1. Panel **c** shows a state of vortex matter which occurs above

the vortex “ionization” temperature. The co-centricity of vortices is strongly reduced, see Fig. 1.

This is also reflected in the subtle difference between protonic (red) and electronic (blue) structure functions in passing from panel **b** to panel **c**. *While the structure function of the electronic vortices essentially is unaffected by passing through the temperature T_C , the structure function for the protonic vortices is distinctly further isotropized.* Namely, the relative increase of the uniform background for the structure function of protonic vortices is a manifestation of the fact that protonic vortices suffer a vortex-loop proliferation transition inside the metallic vortex liquid phase. We have argued above that this transition belongs to the $3Dxy$ universality class, i.e. the same universality class as the superfluid-to-normal fluid transition in liquid helium ⁴He¹⁷.

Concluding, we report the observation, in a numerical experiment, of a novel dissipationless quantum state of matter, namely the *metallic superfluid*. Such a state might be realized in hydrogen or deuterium, if those systems were to take up a projected low-temperature *liquid metallic state* at an extreme pressure. The highest pressure obtained to date appears to be around 320 GPa¹⁸. However, recent breakthroughs in artificial ultrahard diamond synthesis technology¹³ represent sig-




FIG. 2: (Color online) Snapshots of the states of vortex matter and momentum space structure functions generated from MC simulations, taken at three different temperatures: $T = 0.50$ ($T < T_M$), $T = 0.72$ ($T_M < T < T_C$), and $T = 0.86$ ($T > T_C$). The snapshots are extracted from a small segment ($15 \times 15 \times 15$) of the vortex system. Each thick frame contains a sideview and a topview of the vortices, as well as the protonic structure function $S^{(1)}(\mathbf{k}_\perp)$ (red) and the electronic structure function $S^{(2)}(\mathbf{k}_\perp)$ (blue). Panel **a**: For $T < T_M$ the vortices are arranged in a co-centered lattice. Electronic (thick blue) and protonic (thin red) vortices only perform small excursions from each other. Both structure functions exhibit a sixfold Bragg pattern characteristic of a vortex lattice. Panel **b**: For $T_M < T < T_C$ the composite vortex lattice has melted as illustrated by the ring pattern in the structure functions. The electronic and protonic vortices perform stronger excursions from each other, but essentially remain co-centered. This is the superfluid metallic phase in which co-directed currents of protonic and electronic Cooper-pairs can propagate without dissipation. Panel **c**: For $T > T_C$ the superfluidity is lost and the electronic and protonic vortices are no longer co-centered. The proliferation of the protonic vortices is reflected in the increase in the uniform background of the protonic structure function.

nificant progress towards achieving extreme pressures in diamond anvil cells. Thus, a metallic superfluid might be the next “super” state of matter to be realized in the laboratory.

This work was funded by the NTNU through the Norwegian High Performance Computing Program, by the

Research Council of Norway, Grant Nos. 157798/432, 158518/431, and 158547/431 (NANOMAT), by STINT and the Swedish Research Council, and NSF grant DMR-0302347. We thank N. W. Ashcroft for numerous discussions.

¹ A. A. Abrikosov, Sov. Phys. JETP **5**, 1174 (1957).

² Z. Tesanovic, Phys. Rev. **B 59**, 6449 (1999).

³ A. K. Nguyen and A. Sudbø, Phys. Rev., **B 60**, 15307 (1999).

⁴ K. Fossheim and A. Sudbø, *Superconductivity: Physics and Applications*, John Wiley & Sons, London (2004).

⁵ E. Babaev, A. Sudbø, and N. W. Ashcroft, Nature **431** 666 (2004).

⁶ J. Jaffe and N. W. Ashcroft, Phys. Rev. B **23**, R6176 (1981); *ibid*, **B 27**, R5852 (1983); K. Mouloupoulos and N. W. Ashcroft, Phys. Rev. Lett., **66**, 2915 (1991); N. W. Ashcroft, J. Phys. A **12**, A129 (2000); *ibid* A **36** 6137 (2003).

⁷ E. Babaev, L. D. Faddeev, and A. J. Niemi, Phys. Rev. B **65**, 100512(R) (2002).

⁸ E. Babaev, Phys. Rev. Lett. **89**, 067001 (2002).

⁹ J. Smiseth, E. Smørgrav, and A. Sudbø, Phys. Rev. Lett. **93**, 077002 (2004).

¹⁰ E. Smørgrav, J. Smiseth, E. Babaev, and A. Sudbø, Phys. Rev. Lett. **94**, 096401 (2005); J. Smiseth, E. Smørgrav, E. Babaev, and A. Sudbø, Phys. Rev. B **71**, 214509 (2005).

¹¹ T. Senthil *et al.*, Science **303**, 1490 (2004).

¹² S. A. Bonev, E. Schwegler, T. Ogitsu, and G. Galli, Nature **431**, 669 (2004).

¹³ For a recent press release, see: Very Large Diamonds Produced Very Fast, Carnegie Institution of Washington, May

- 16, 2005.
- ¹⁴ E. Gregoryanz *et al.*, Phys. Rev. Lett., **94**, 185502 (2005).
- ¹⁵ T. Guillot, Physics Today **57**, 63 (2004).
- ¹⁶ E. W. Kim and M. H. W. Chan, Nature **427**, 225 (2004).
- ¹⁷ H. Kleinert, *Gauge Fields in Condensed Matter Physics*, (World Scientific, Singapore, 1989), Vol. 1.
- ¹⁸ P. Loubeyre, F. Ocelli, and R. Le Toullec, Nature, **416**, 613 (2002).